

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
30 October 2003 (30.10.2003)

PCT

(10) International Publication Number  
**WO 03/089637 A1**

(51) International Patent Classification<sup>7</sup>: C12N 9/12, 15/03

(74) Agent: BANFORD, P., C.; Marks & Clerk, 43 Park Place,  
Leeds LS1 2RY (GB).

(21) International Application Number: PCT/GB03/01623

(22) International Filing Date: 15 April 2003 (15.04.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0208768.2 17 April 2002 (17.04.2002) GB

(71) Applicant (for all designated States except US): **THE UNIVERSITY OF NEWCASTLE** [GB/GB]; 6 Kensington Terrace, Jesmond, Newcastle upon Tyne NE1 7RU (GB).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and


(75) Inventors/Applicants (for US only): **CONNOLLY, Bernard** [GB/GB]; School of Cell and Molecular Biosciences, University of Newcastle, Newcastle upon Tyne NE2 4HH (GB). **FOGG, Mark** [GB/GB]; Structural Biology Lab, Department of Chemistry, University of York, York YO10 5YW (GB). **PEARL, Laurence** [GB/GB]; Centre for Structural Biology, Institute of Cancer Research, Chester Beatty Laboratories, 237 Fulham Road, London SW3 6JB (GB).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: MUTATION OF DNA POLYMERASES FROM ARCHAEOBACTERIA



(57) Abstract: The present invention relates to a variant archaeal DNA polymerase having a modified amino acid sequence of a wild-type amino acid sequence, the modified sequence being in the amino-terminal amino acids that comprise a uracil-binding pocket in the wild-type polymerase whereby the variant polymerase has reduced affinity for uracil than the wild-type polymerase. Such variant polymerases may be usefully employed in biological assay systems such as the polymerase chain reaction.

**BEST AVAILABLE COPY**

WO 03/089637 A1

## MUTATION OF DNA POLYMERASES FROM ARCHAEOBACTERIA

The present invention relates to archaeal DNA polymerase variants and their use in the amplification of DNA.

Polymerase chain reaction (PCR) is a method whereby a sequence of DNA may be selectively amplified to produce a large sample that may be readily analyzed. A solution containing the DNA to be amplified, together with free bases, a polymerase enzyme and primers that bind to opposite ends of the two strands of the DNA segment to be replicated, is heated to break the bonds between the strands of DNA. When the solution cools, the primers bind to the separated strands and the polymerase builds a new strand by joining free bases to the primers thereby producing a new strand that is restricted solely to the desired segment. PCR enables billions of copies of a small piece of DNA to be produced in several hours.

Heat stable polymerases are required for this process and one of the most commonly used is Taq DNA polymerase from *Thermus aquaticus*. However, this enzyme does not possess a 3'-5' exonuclease III function, commonly referred to as "proofreading activity". This function removes bases that are mismatched at the 3' end of a primer-template duplex. The inability of Taq DNA polymerase to carry out this function results in it being prone to base incorporation errors.

Archaeal DNA polymerases are thermally stable and demonstrate proofreading activity. However, native archaeal DNA polymerases are inhibited by deoxyuracil. Archaeal DNA polymerases have a "read-ahead" function specifically for uracil. This template

checking activity scans the template ahead of the replication fork for the presence of uracil and stalls polymerisation when uracil is encountered. Thus, the presence of deoxyuracil in DNA causes amplification to be stalled when using native archaeal DNA polymerases. This is a serious setback since the repetitive heating and cooling cycles of a DNA sample being amplified by PCR results in partial, thermally induced deamination of dCTP (a component incorporated into newly amplified DNA) to dUTP (which can be incorporated into DNA) and deamination of deoxycytidine in the DNA to deoxyuracil. This can result in the native archaeal DNA polymerases being unsuitable for PCRs, in particular those concerned with the prevention of "carry-over contamination" where PCR is carried out with dUTP rather than dTTP.

It is an object of the present invention to provide modified archaeal DNA polymerases that do not have the disadvantage of being inhibited by deoxyuracil and are particularly useful in polymerase chain reactions.

Accordingly, a first aspect of the present invention provides a variant archaeal DNA polymerase having a modified amino acid sequence of a wild-type amino acid sequence, the modified sequence being in the amino-terminal amino acids that comprise a uracil-binding pocket in the wild-type polymerase whereby the variant polymerase has reduced affinity for uracil compared to wild-type archaeal DNA polymerases.

The present invention is based upon research (see the Examples) conducted by the inventors that has identified a uracil-binding pocket in archaeal DNA polymerases. They

realised that this pocket may be altered to provide variant polymerases according to the invention that may be beneficially used as described herein.

The variant archaeal DNA polymerase may be a modification of an archaeal family B DNA polymerase. For instance the variant may be derived from any one of the fourteen archaeal family B DNA polymerase shown in Figure 1. For instance the variant may be derived from the polymerases found in *Pyrococcus furiosus* (Pfu-Pol), *Thermococcus gorgonarius* (Tgo-Pol), *Thermococcus litoralis* (Tli-Pol), *Thermococcus* sp. 9°N-7 (9°N-7-Pol), *Desulfurococcus* strain Tok (DTok-Pol), *Pyrobaculum islandicum* (Pis-Pol), *Archaeoglobus fulgidus* (Afu-Pol), *Sulfolobus acidocaldarius* (Sac-Pol), *Sulfurisphaera ohwakuensis* (Soh-Pol), *Sulfolobus solfataricus* (Sso-Pol), *Pyrodictium occultum* (Poc-Pol) or *Aeropyrum pernix* (Ape-Pol). It will be appreciated that the variant could also be derived from any other archaeal family B DNA polymerase.

It is preferred that the variant is derived from *Pyrococcus furiosus* (Pfu-Pol). Pfu-Pol used by the inventors has the following amino acid sequence:

```
MAILDVDYITEEGKPVIRLFKKENGKFKIEHDRTFRPYIYALLRDDSKIEEVKKITGERHKGKIVRIVDVEK
VEKKFLGKPITVWKLYLEHPQDVPTIREKVREHPAVVDIFEYDIPFAKRYLIDKGLIPMEGEEELKILAF
DIETLYHEGEEFGKGPIIMISYADENEAKVITWKNIDLPYVEVSSEREMIKRFLRIIREKDPDIIVTYN
GDSFDFPYLAKRAEKLGIKLTIGRDGSEPKMQRIGDMTAVEVKGRIFDLYHVITRTINLPTYTLEAVYE
AIFGKPKKEKVYADEIAKAWESGENLERVAKYSMEDAKATYELGKEFLPMEIQLSRLVGQPLWDVSRSTG
NLVEWFLLRKAYERNEVAPNKPSEEEYQRRRLRESYTGGFVKEPEKGLWENIVYLDLFRALYPSIIITHNVS
PDTLNLEGCKNYDIAPQVGHKFKCDIPGFIPSLLGHLLEERQKIKTMMKETQDPIEKILLDYRQKAIKLL
ANSFYGYGYAKARWYCKEACSVTAWGRKYIELVWKELEEKFGFKVLYIDTDGLYATIPGGESEEEKKK
ALEFVKYINSLPGLLELEYEGFYKRGFFVTKKRYAVIDEEGKVITRGLEIVRRDWSEIAKETQARVLET
ILKHGDFVEEAVRIVKEVIQKLANYEIPPEKLAIYEQITRPLHEYKAIGPHVAVAKKLAAGVKIKPGMVI
GYIVLRGDPISNRAILAEEDPKKHKYDAEYYIENQVLPVLRILEGFGYRKEDLRYQKTRQVGLTSWL
NIKKK
```

(SEQ ID NO.1)

The Pfu Pol used by the inventors contains an extra A at position 2. This extra amino acid was incorporated because it improves protein expression without affecting the properties of the enzyme. The true wild type Pfu Pol begins MILDVDY. The sequence of the true wild type is:

```
MILDVDYITEEGKPVIRLFKKENGKFKIEHRTFRPYIYALLRDDSKIEEVKKITGERHKGKIVRIVDVEK
VEKKFLGKPITVWKLYLEHPQDVPTIREKVREHPAVVDIFEYDIPFAKRYLIDKGLIPMEGEEELKILAF
DIETLYHEGEEFGKGPIIMISYADENEAKVITWKNIDLPYVEVVSSEREMIKRFLRIIREKDPDIIVTYN
GDSFDFPYLAKRAEKLGIKLTIGRDGSEPKMQRIGDMTAVEVKGRIHFDLYHVITRTINLPTYTLEAVYE
AIFGKPKEKVYADEIAKAWESGENLERVAKYSMEDAKATYELGKEFLPMEIQLSRLVGQPLWDVSRSTG
NLVEWFLLRKAYERNEVAPNKPSEEEYQRRRLRESYTGGFVKPEKGLWENIVYLDFRALYPSIIITHNVS
PDTLNLEGCKNYDIAPQVGHKFKDIPGFIPSLGLHLEERQKIKTKMKETQDPIEKILLDYRQKAIKLL
ANSFYGYGYAKARWYCKECAESVTAWGRKYIELVWKELEEKFGFKVLYIDTDGLYATIPGGESEEEKK
ALEFVKYINSLPGLLELEYEGFYKRGFFVTCKRYAVIDEKGKIVITRGLIVRRDWSEIAKETQARVLET
ILKHGDEV EAVRIVKEVIQKLANYEIPPEKLAIYEQITRPLHEYKAIGPHVAVAKKLAAGVKIKPGMVI
GYIVLRGDPISNRAILAEYDPPKKHKYDAEYYIENQVLPVAVLRILEGFGYRKEDLRYQKTRQVGLTSWL
NIKKS
```

(SEQ ID NO.2)

Accordingly, preferred mutants according to the invention begin MAILDVDY or MILDVDY.

The inventors have found (see Example 1) that a uracil-binding pocket of the wild type polymerase forms part of the ssDNA template binding cleft of the polymerase (the so call cleft T). Furthermore the uracil-binding pocket comprises amino acids from two conserved regions of the polymerases: Region A and Region B separated by an unconserved region. In the archaeal polymerase from *Pyrococcus furiosus* (Pfu-Pol) Region A is formed by the amino acids 1-40 and Region B by amino acids 78-130. Highly conserved residues in these two regions form the highly ordered uracil-binding pocket. Other archaea have similar regions A & B in their respective polymerases as illustrated in Figure 1. Preferably, one or more of the amino acids in Regions A and/or B are altered to form the variant archaeal DNA polymerase.

Figure 1 illustrates a sequence alignment of the N -terminal domains of various archaeal polymerases. In figure 1 amino acids designated (1) have 90% or greater identity, (2) indicates 80- 90% identity and (3) 60-80% identity. The two highly conserved regions that form the uracil binding pocket are:

Region A, amino acids 1-40 in Pfu-Pol (and corresponding regions in the other polymerases); and

Region B, amino acids 78-131 in Pfu-Pol (and corresponding regions in the other polymerases).

It is preferred that the variant is formed by alteration of one of the amino acids block shaded (1, 2 or 3) in Figure 1. For example the alteration may be in the motif: E - - I - F/Y- - -Y- -D.

The alteration may consist of a substitution, deletion or addition. One of the invariant residues may be altered or other residues in Regions A and/or B that affect the conformation of the uracil-binding pocket.

The inventors believe that residues 7, 36, 37, 90-97 and 112 - 119 in Pfu-Pol are particularly important for uracil binding. Preferably, at least one of these residues is altered to effect the conformation of the pocket and thereby reduce its uracil-binding ability. More preferably, the mutation in Pfu-Pol consists of a change in the amino acids Y7, Y37, V93, I114 or P115. More preferably, the change consists of Y7A, Y37A, V93Q, V93R, I114R, I114Q or P115A. A most preferred Pfu-Pol mutation is V93Q.

Examples of preferred Pfu-Pol variants have the following amino acid sequences:

(a) Pfu pol Y7A (Y8A)

MAILDVDVAITEEGKPVIRLFKKENGKFKIEHRTFRPYIYALLRDDSKEEVKKITGERHGKIVRIVDVEK  
VEKKFLGKPITVWKLYLEHPQDVPTIREKVREHPAVVDIFEYDIPFAKRYLIDKGLIPMEGEEELKILAF  
DIETLYHEGEEFGKGPIIMISYADENEAKVITWKNIDLPYVEVVSSEMERIKRFLRIIREKDPDIIVTYN  
GDSFDFPYLAKRAEKLGIKLTIGRDGSEPKMQRIGDMTAVEVKGRIFDLYHVITRTINLPTYTLEAVYE  
AIFGKPKKEKVYADEIAKAWESGENLERVAKYSMEDAKATYELGKEFLPMEIQLSRLVGQPLWDVSRSTG  
NLVEWFLLRKAYERNEVAPNKPSEEEYQRRRLRESYTGGFVKEPEKGLWENIVYLDLFRALYPSIIITHNVS  
PDTLNLEGCKNYDIAPQVGHKFCCKDIPGFIPSLLGHLLEERQKIKTKMKETQDPIEKILLDYRQKAIKLL  
ANSFYGYGYAKARWYCKECAESVTAWGRKYIELVWKELEEKFGFKVLYIDTDGLYATIPGGESEEEIKKK  
ALEFVKYINSLPGLLELEYEGFYKRGFFVTKKRYAVIDEKGKIVITRGLIVRRDWSEIAKETQARVLET  
ILKHGDVEEAVRIVKEVIQKLANYEIPPEKLAIYEQITRPLHEYKAIGPHVAVAKKLAAGVKIKPGMVI  
GYIVLRGDPISNRAILAEYDPPKKHKYDAEYYIENQVLPVLRILEGFGYRKEDLRYQKTRQVGLTSLW  
NIKKK

(SEQ ID NO.3)

(b) Pfu pol Y37A (Y38A)

MAILVDYITEEGKPVIRLFKKENGKFKIEHRTFRPAIYALLRDDSKEEVKKITGERHGKIVRIVDVEK  
VEKKFLGKPITVWKLYLEHPQDVPTIREKVREHPAVVDIFEYDIPFAKRYLIDKGLIPMEGEEELKILAF  
DIETLYHEGEEFGKGPIIMISYADENEAKVITWKNIDLPYVEVVSSEMERIKRFLRIIREKDPDIIVTYN  
GDSFDFPYLAKRAEKLGIKLTIGRDGSEPKMQRIGDMTAVEVKGRIFDLYHVITRTINLPTYTLEAVYE  
AIFGKPKKEKVYADEIAKAWESGENLERVAKYSMEDAKATYELGKEFLPMEIQLSRLVGQPLWDVSRSTG  
NLVEWFLLRKAYERNEVAPNKPSEEEYQRRRLRESYTGGFVKEPEKGLWENIVYLDLFRALYPSIIITHNVS  
PDTLNLEGCKNYDIAPQVGHKFCCKDIPGFIPSLLGHLLEERQKIKTKMKETQDPIEKILLDYRQKAIKLL  
ANSFYGYGYAKARWYCKECAESVTAWGRKYIELVWKELEEKFGFKVLYIDTDGLYATIPGGESEEEIKKK  
ALEFVKYINSLPGLLELEYEGFYKRGFFVTKKRYAVIDEKGKIVITRGLIVRRDWSEIAKETQARVLET  
ILKHGDVEEAVRIVKEVIQKLANYEIPPEKLAIYEQITRPLHEYKAIGPHVAVAKKLAAGVKIKPGMVI  
GYIVLRGDPISNRAILAEYDPPKKHKYDAEYYIENQVLPVLRILEGFGYRKEDLRYQKTRQVGLTSLW  
NIKKK

(SEQ ID NO.4)

(c) Pfu pol V93Q (V94Q)

MAILVDYITEEGKPVIRLFKKENGKFKIEHRTFRPYIYALLRDDSKEEVKKITGERHGKIVRIVDVEK  
VEKKFLGKPITVWKLYLEHPQDVPTIREKVREHPAVVDIFEYDIPFAKRYLIDKGLIPMEGEEELKILAF  
DIETLYHEGEEFGKGPIIMISYADENEAKVITWKNIDLPYVEVVSSEMERIKRFLRIIREKDPDIIVTYN  
GDSFDFPYLAKRAEKLGIKLTIGRDGSEPKMQRIGDMTAVEVKGRIFDLYHVITRTINLPTYTLEAVYE  
AIFGKPKKEKVYADEIAKAWESGENLERVAKYSMEDAKATYELGKEFLPMEIQLSRLVGQPLWDVSRSTG  
NLVEWFLLRKAYERNEVAPNKPSEEEYQRRRLRESYTGGFVKEPEKGLWENIVYLDLFRALYPSIIITHNVS  
PDTLNLEGCKNYDIAPQVGHKFCCKDIPGFIPSLLGHLLEERQKIKTKMKETQDPIEKILLDYRQKAIKLL  
ANSFYGYGYAKARWYCKECAESVTAWGRKYIELVWKELEEKFGFKVLYIDTDGLYATIPGGESEEEIKKK  
ALEFVKYINSLPGLLELEYEGFYKRGFFVTKKRYAVIDEKGKIVITRGLIVRRDWSEIAKETQARVLET  
ILKHGDVEEAVRIVKEVIQKLANYEIPPEKLAIYEQITRPLHEYKAIGPHVAVAKKLAAGVKIKPGMVI  
GYIVLRGDPISNRAILAEYDPPKKHKYDAEYYIENQVLPVLRILEGFGYRKEDLRYQKTRQVGLTSLW  
NIKKK

(SEQ ID NO.5)

## (d) Pfu pol V93R (V94R)

MAILDVDYITEEGKPVIRLFKKENGKFKIEHRTFRPYIYALLRDDSKEEVKKITGERHGKIVRIVDVEK  
VEKKFLGKPITVWKLYLEHPQDRPTIREKVREHPAVVDIFEYDIPFAKRYLIDKGLIPMEGEEELKILAF  
DIETLYHEGEEFGKGPIIMISYADENEAKVITWKNIDLPHYVEVVSSEREMIKRFLRIIREKDPDIIVTYN  
GDSFDFPYLAKRAEKLGIKLTIGRDGSEPKMQRIGDMTAVEVKGRIHFDLYHVITRTINLPTYTLEAVYE  
AIFGKPKKVVYADEIAKAWESGENLERVAKYSMEDAKATYELGKEFLPMEIQLSRLVGQPLWDVSRSTG  
NLVEWFLLRKAYERNEVAPNKPSEEEYQRRRESYTGGFVKEPEKGLWENIVYLDLFRALYPSIIITHNVS  
PDTLNLEGCKNYDIAPOVGHKFCKDIPGFIPSLLGHLLEERQKIKTKMKETQDPIEKILLDYRQKAIKLL  
ANSFYGYGYAKARWYCKECAESVTAWGRKYIELVWKELEEKFGFKVLYIDTDGLYATIPGGESEEEIKKK  
ALEFVKYINSKLPGLLELEYEGFYKRGFFVTKKRYAVIDEEGKVITRGLIVRRDWSEIAKETQARVLET  
ILKHGDVEEAVRIVKEVIQKLANYEIPPEKLAIYEQITRPLHEYKAIGPHVAVAKKLAAGVKIKPGMVI  
GYIVLRGDPISNRILAEEDPKKHKYDAEYYIENQVLPVLRILEGFGYRKEDLRYQKTRQVGLTSWL  
NIKKK

(SEQ ID NO.6)

## (e) Pfu pol I114R (I115R)

MAILDVDAITTEEGKPVIRLFKKENGKFKIEHRTFRPYIYALLRDDSKEEVKKITGERHGKIVRIVDVEK  
VEKKFLGKPITVWKLYLEHPQDVPTIREKVREHPAVVDIFEYDPPFAKRYLIDKGLIPMEGEEELKILAF  
DIETLYHEGEEFGKGPIIMISYADENEAKVITWKNIDLPHYVEVVSSEREMIKRFLRIIREKDPDIIVTYN  
GDSFDFPYLAKRAEKLGIKLTIGRDGSEPKMQRIGDMTAVEVKGRIHFDLYHVITRTINLPTYTLEAVYE  
AIFGKPKKVVYADEIAKAWESGENLERVAKYSMEDAKATYELGKEFLPMEIQLSRLVGQPLWDVSRSTG  
NLVEWFLLRKAYERNEVAPNKPSEEEYQRRRESYTGGFVKEPEKGLWENIVYLDLFRALYPSIIITHNVS  
PDTLNLEGCKNYDIAPOVGHKFCKDIPGFIPSLLGHLLEERQKIKTKMKETQDPIEKILLDYRQKAIKLL  
ANSFYGYGYAKARWYCKECAESVTAWGRKYIELVWKELEEKFGFKVLYIDTDGLYATIPGGESEEEIKKK  
ALEFVKYINSKLPGLLELEYEGFYKRGFFVTKKRYAVIDEEGKVITRGLIVRRDWSEIAKETQARVLET  
ILKHGDVEEAVRIVKEVIQKLANYEIPPEKLAIYEQITRPLHEYKAIGPHVAVAKKLAAGVKIKPGMVI  
GYIVLRGDPISNRILAEEDPKKHKYDAEYYIENQVLPVLRILEGFGYRKEDLRYQKTRQVGLTSWL  
NIKKK

(SEQ ID NO.7)

## (f) Pfu pol I114Q (I115Q)

MAILDVDYITEEGKPVIRLFKKENGKFKIEHRTFRPAIYALLRDDSKEEVKKITGERHGKIVRIVDVEK  
VEKKFLGKPITVWKLYLEHPQDVPTIREKVREHPAVVDIFEYDQPPFAKRYLIDKGLIPMEGEEELKILAF  
DIETLYHEGEEFGKGPIIMISYADENEAKVITWKNIDLPHYVEVVSSEREMIKRFLRIIREKDPDIIVTYN  
GDSFDFPYLAKRAEKLGIKLTIGRDGSEPKMQRIGDMTAVEVKGRIHFDLYHVITRTINLPTYTLEAVYE  
AIFGKPKKVVYADEIAKAWESGENLERVAKYSMEDAKATYELGKEFLPMEIQLSRLVGQPLWDVSRSTG  
NLVEWFLLRKAYERNEVAPNKPSEEEYQRRRESYTGGFVKEPEKGLWENIVYLDLFRALYPSIIITHNVS  
PDTLNLEGCKNYDIAPOVGHKFCKDIPGFIPSLLGHLLEERQKIKTKMKETQDPIEKILLDYRQKAIKLL  
ANSFYGYGYAKARWYCKECAESVTAWGRKYIELVWKELEEKFGFKVLYIDTDGLYATIPGGESEEEIKKK  
ALEFVKYINSKLPGLLELEYEGFYKRGFFVTKKRYAVIDEEGKVITRGLIVRRDWSEIAKETQARVLET  
ILKHGDVEEAVRIVKEVIQKLANYEIPPEKLAIYEQITRPLHEYKAIGPHVAVAKKLAAGVKIKPGMVI  
GYIVLRGDPISNRILAEEDPKKHKYDAEYYIENQVLPVLRILEGFGYRKEDLRYQKTRQVGLTSWL  
NIKKK

(SEQ ID NO.8)



It will be appreciated that the preferred mutants listed above comprise an alanine (A) insertion at position 2 which is not found in the wild type. Accordingly such mutants may be designated Y8A, Y38A, V94Q, V94R, I115R and I115Q mutants of the MAILDVDY form of Pfu Pol and correspond to Y7A, Y37A, V93Q, V93R, I114R and I114Q mutants of the true wild-type (MILDVDY). Y7A, Y37A, V93Q, V93R, I114R and I114Q mutants of the true wild-type (MILDVDY) are also preferred mutants according to the invention.

It will be appreciated that equivalent residues in other archaeal polymerases may be mutated (see figure 1). For instance Y7A, Y37A, V93Q, V93R, I114R, I114Q and P115Δ remain preferred mutants in Tgo-pol, DTok-pol and 9°N-7-pol.

The present invention also provides nucleic acids encoding an archaeal DNA polymerase variant as defined above.

Examples of preferred mutant genes have the following DNA sequences:

**(a) Pfu pol Y7A (Y8A)**

```
ATGGCTATCCTGGACGTTGACGCCATCACCGAAGAAGGTAAGCCGGTTATCCGTCTGTTCAAAAAGAAAACGGT
AAATTCAAAATCGAACACGACCGTACCTTCCGTCCGTACATCTACGCTCTGCTGCGTGACGACTCTAAAATCGAA
GAAGTTAAAAAATCACCGGTGAACGTCATGGAAAGATTGTGAGAATTGTTGATGTAGAGAAGGTTGAGAAAAAG
TTTCTCGGCAAGCCTATTACCGTGTGGAACTTTATTTGGAACATCCCCAAGATGTTCCCACTATTAGAGAAAA
GTTAGAGAACATCCAGCAGTTGTGGACATCTTCGAATACGATATTCCATTTGCAAAGAGATACCTCATCGACAAA
GGCCTAATACCAATGGAGGGGGAAGAAGAGCTAAAGATTCTTGCCCTTCGATATAGAAACCCTCTATCACGAAGGA
GAAGAGTTTGGAAAAGGCCCAATTATAATGATTAGTTATGCAGATGAAAATGAAGCAAAGGTGATTACTTGGAAA
AACATAGATCTTCCATACGTTGAGGTTGTATCAAGCGAGAGAGAGATGATAAAGAGATTTCTCAGGATTATCAGG
GAGAAGGATCCTGACATTATAGTTACTTTATAATGGAGACTCATTCGACTTCCCATATTTAGCGAAAAGGGCAGAA
AACTTGGGATTAAATTAACCATTTGGAAGAGATGGAAGCGAGCCCAAGATGCAGAGAATAGGCGATATGACGGCT
GTAGAAGTCAAGGGAAGAATACATTTGCACTTGTATCATGTAATAACAAGGACAATAAATCTCCCAACATACACA
CTAGAGGCTGTATATGAAGCAATTTTGGAAAGCCAAAGGAGAAGGTATACGCCGACGAGATAGCAAAAGCCTGG
GAAAGTGGAGAGAACCTTGAGAGAGTTGCCAAATACTCGATGGAAGATGCAAAGGCAACTTATGAACTCGGGAAA
GAATTCCTTCCAATGGAAATTCAGCTTTCAAGATTAGTTGGACAACCTTTATGGGATGTTTCAAGGTCAAGCACA
GGGAACCTTGTAGAGTGGTTCTTACTTAGGAAAAGCCTACGAAAGAAACGAAGTAGCTCCAAACAAGCCAAGTGAA
GAGGAGTATCAAGAAGGCTCAGGGAGAGCTACACAGGTGGATTTCGTTAAAGAGCCAGAAAAGGGGTTGTGGGAA
AACATAGTATACCTAGATTTTAGAGCCCTATATCCCTCGATTATAATTACCCACAATGTTTCTCCCGATACTCTA
```

AATCTTGAGGGGATGCAAGAACTATGATATCGCTCCTCAAGTAGGCCACAAGTTCTGCAAGGACATCCCTGGTTTT  
ATACCAAGTCTCTTGGGACATTTGTTAGAGGAAAGACAAAAGATTAAGACAAAATGAAGGAACTCAAGATCCT  
ATAGAAAAAATACTCCTTGACTATAGACAAAAGCGATAAAACTCTTAGCAAATTTCTTCTACGGATATTATGGC  
TATGCAAAAGCAAGATGGTACTGTAAGGAGTGTGCTGAGAGCGTTACTGCCTGGGGAAGAAAGTACATCGAGTTA  
GTATGGAAGGAGCTCGAAGAAAAGTTTGGATTTAAAGTCTCTACATTGACACTGATGGTCTCTATGCAACTATC  
CCAGGAGGAGAAAGTGAGGAAATAAAGAAAAAGGCTCTAGAATTTGTAAAATACATAAATTCAAAGCTCCCTGGA  
CTGCTAGAGCTTGAATATGAAGGGTTTTATAAGAGGGGATTTCTTCTGTTACGAAGAAGAGGTATGCAGTAATAGAT  
GAAGAAGGAAAAGTCATTACTCGTGGTTTTAGAGATAGTTAGGAGAGATTGGAGTGAAATTGCAAAAGAACTCAA  
GCTAGAGTTTTGGAGACAATACTAAAACACGGAGATGTTGAAGAAGCTGTGAGAATAGTAAAGAAGTAATACAA  
AAGCTTGCCAATTATGAAATTCCACCAGAGAAGCTCGCAATATATGAGCAGATAACAAGACCATTACATGAGTAT  
AAGGCGATAGGTCCTCACGTGTTGTCGAAAGAACTGCTGCTTAAAGGAGTTAAAATAAAGCCAGGAATGGTA  
ATTGGATACATAGTACTTAGAGGCGATGGTCCAATTAGCAATAGGGCAATTCTAGCTGAGGAATACGATCCCAAA  
AAGCACAAGTATGACGCAGAATATTACATTGAGAACCAGGTTCTTCCAGCGGTACTTAGGATATTGGAGGGATTT  
GGATACAGAAAGGAAGACCTCAGATACCAAAAGACAAGACAAGTCGGCCTAACTTCCTGGCTTAACATTAAAAAA  
TCC

(SEQ ID NO.9)

**(b) Pfu pol V93Q (V94Q)**

ATGGCTATCCTGGACGTTGACTACATCACCGAAGAAGGTAAGCCGGTTATCCGTCTGTTCAAAAAAGAAAACGGT  
AAATTCAAAATCGAACACGACCGTACCTTCCGTCCGTACATCTACGCTCTGCTGCGTGACGACTCTAAAATCGAA  
GAAGTTAAAAAATCACCGGTGAACGTCATGGAAGATTGTGAGAATTGTTGATGTAGAGAAGGTTGAGAAAAAG  
TTTCTCGGCAAGCCTATTACCGTGTGGAACTTTATTTGGAACATCCCCAAGATCAGCCCACTATTAGAGAAAA  
GTTAGAGAACATCCAGCAGTTGTGGACATCTTCGAATACGATATTCCATTGCAAGAGATACCTCATCGACAAA  
GGCCTAATACCAATGGAGGGGGAAGAAGAGCTAAAGATTCTTGCCCTTCGATATAGAAAACCTCTATCAGGAAGGA  
GAAGAGTTTGGAAAAGGCCCAATTATAATGATTAGTTATGCAGATGAAAATGAAGCAAAGGTGATTACTTGGAAA  
AACATAGATCTTCCATACGTTGAGGTTGTATCAAGCGAGAGAGAGATGATAAAGAGATTCTCAGGATTATCAGG  
GAGAAGGATCCTGACATTATAGTTACTTATAATGGAGACTCATTGCACTTCCCATATTTAGCGAAAAGGGCAGAA  
AACTTGGGATTAAATTAACCATTGGAAGAGATGGAAGCGAGCCCAAGATGCAGAGAATAGGCGATATGACGGCT  
GTAGAAGTCAAGGGAAGAATACATTTGCACTTGTATCATGTAATAACAAGGACAATAAATCTCCCAACATACACA  
CTAGAGGCTGTATATGAAGCAATTTTTGGAAAGCCAAAGGAGAAGGTATACGCCGACGAGATAGCAAAAGCCTGG  
GAAAGTGGAGAGAACCTTGAGAGAGTTGCCAAATACTCGATGGAAGATGCAAAGGCAACTTATGAACTCGGGAAA  
GAATTCCTTCCAATGGAAATTCAGCTTTCAAGATTAGTTGGACAACCTTTATGGGATGTTTCAAGGTCAAGCACA  
GGGAACCTTGTAGAGTGGTTCTTACTTAGGAAAGCTACGAAAGAAACGAAGTAGCTCCAAACAGCCAAGTAA  
GAGGAGTATCAAAGAAAGGCTCAGGGAGAGCTACACAGGTGGATTTCGTTAAAGAGCCAGAAAAGGGGTTGTGGGAA  
AACATAGTATACCTAGATTTTAGAGCCCTATATCCCTCGATTATAATTACCCACAATGTTTCTCCCGATACTCTA  
AATCTTGAGGGATGCAAGAACTATGATATCGCTCCTCAAGTAGGCCACAAGTTCTGCAAGGACATCCCTGGTTTT  
ATACCAAGTCTCTTGGGACATTTGTTAGAGGAAAGACAAAAGATTAAGACAAAATGAAGGAACTCAAGATCCT  
ATAGAAAAAATACTCCTTGACTATAGACAAAAGCGATAAAACTCTTAGCAAATTTCTTCTACGGATATTATGGC  
TATGCAAAAGCAAGATGGTACTGTAAGGAGTGTGCTGAGAGCGTTACTGCCTGGGGAAGAAAGTACATCGAGTTA  
GTATGGAAGGAGCTCGAAGAAAAGTTTGGATTTAAAGTCTCTACATTGACACTGATGGTCTCTATGCAACTATC  
CCAGGAGGAGAAAAGTGAGGAAATAAAGAAAAAGGCTCTAGAATTTGTAAAATACATAAATTCAAAGCTCCCTGGA  
CTGCTAGAGCTTGAATATGAAGGGTTTTATAAGAGGGGATTCTTCTGTTACGAAGAAGAGGTATGCAGTAATAGAT  
GAAGAAGGAAAAGTCATTACTCGTGGTTTTAGAGATAGTTAGGAGAGATTGGAGTGAAATTGCAAAAGAACTCAA  
GCTAGAGTTTTTGGAGACAATACTAAAACACGGAGATGTTGAAGAAGCTGTGAGAATAGTAAAAGAAGTAATACAA  
AAGCTTGCCAATTATGAAATTCCACCAGAGAAGCTCGCAATATATGAGCAGATAACAAGACCATTACATGAGTAT  
AAGGCGATAGGTCCTCACGTAGCTGTTGCAAGAACTAGCTGCTTAAAGGAGTTAAAATAAAGCCAGGAATGGTA  
ATTGGATACATAGTACTTAGAGGCGATGGTCCAATTAGCAATAGGGCAATTCTAGCTGAGGAATACGATCCCAAA  
AAGCACAAGTATGACGCAGAATATTACATTGAGAACCAGGTTCTTCCAGCGGTACTTAGGATATTGGAGGGATTT  
GGATACAGAAAGGAAGACCTCAGATACCAAAAGACAAGACAAGTCGGCCTAACTTCCTGGCTTAACATTAAAAAA  
TCC

(SEQ ID NO.10)

## (c) Pfu Pol P115Δ (P116Δ)

ATGGCTATCCTGGACGTTGACTACATCACCGAAGAAGGTAAGCCGGTTATCCGTCTGTTCAAAAAAGAAAAACGGT  
AAATTCAAAATCGAACACGACCGTACCTTCCGTCCGTACATCTACGCTCTGCTGCGTGACGACTCTAAAATCGAA  
GAAGTTAAAAAATCACCGGTGAACGTCATGGAAAGATTGTGAGAATTGTTGATGTAGAGAAGGTTGAGAAAAAG  
TTTCTCGGCAAGCCTATTACCGTGTGGAAACTTTATTTGGAAACATCCCAAGATGTTCCCACTATTAGAGAAAAA  
GTTAGAGAACATCCAGCAGTTGTGGACATCTTCGAATACGATATTTTTGCAAAGAGATACCTCATCGACAAAGGC  
CTAATACCAATGGAGGGGGAAGAAGAGCTAAAGATTCTTGCCTTCGATATAGAAACCCCTCTATCACGAAGGAGAA  
GAGTTTGGAAAAGGCCCAATTATAATGATTAGTTATGCAGATGAAAATGAAGCAAGGTTGATTACTTGGAAAAAC  
ATAGATCTTCCATACGTTGAGGTTGTATCAAGCGAGAGAGAGATGATAAAGAGATTTCTCAGGATTATCAGGGAG  
AAGGATCCTGACATTATAGTTACTTATAATGGAGACTCATTCGACTTCCCATATTTAGCGAAAAAGGCGAGAAAAA  
CTTGGGATTAAATTAACCATTTGGAAGAGATGGAAGCGAGCCCAAGATGCAGAGAATAGGCGATATGACGGCTGTA  
GAAGTCAAGGGAAGAATACATTTGCACTTGTATCATGTAATAACAAGGACAATAAATCTCCCAACATACACACTA  
GAGGCTGTATATGAAGCAATTTTTGGAAAGCCAAAGGAGAAGGTATACGCCGACGAGATAGCAAAAGCCTGGGAA  
AGTGGAGAGAACCCTTGAGAGAGTTGCCAAATACTCGATGGAAGATGCAAAGGCAACTTATGAACTCGGGAAAGAA  
TTCCCTTCCAATGGAAATTCAGCTTTCAAGATTAGTTGGACAACCTTTATGGGATGTTTCAAGGTCAAGCACAGGG  
AACCTTGTAGAGTGGTTCTTACTTAGGAAAGCCTACGAAAGAAACGAAGTAGCTCCAAACAAGCCAAGTGAAGAG  
GAGTATCAAAGAAGGCTCAGGGAGAGCTACACAGGTGGATTTCGTTAAAGAGCCAGAAAAGGGGTTGTGGGAAAAC  
ATAGTATACCTAGATTTTAGAGCCCTATATCCCTCGATTATAATTACCCACAATGTTTCTCCCGATACTCTAAAT  
CTTGAGGGATGCAAGAAGCTATGATATCGCTCCTCAAGTAGGCCACAAGTTCTGCAAGGACATCCCTGGTTTTTATA  
CCAAGTCTCTTGGGACATTTGTTAGAGGAAGCAAAAGGATAAAACTCTTAGCAAATTCTTTCTACGGATATTATGGCTAT  
GAAAAAATACTCCTTGACTATAGACAAAAGCGATAAAACTCTTAGCAAATTCTTTCTACGGATATTATGGCTAT  
GCAAAAGCAAGATGGTACTGTAAGGAGTGTGCTGAGAGCGTTACTGCCTGGGGAAGAAAGTACATCGAGTTAGTA  
TGGAAGGAGCTCGAAGAAAAGTTTGGATTTAAAGTCCTCTACATTGACACTGATGGTCTCTATGCAACTATCCCA  
GGAGGAGAAAAGTGGAGAAATAAAGAAAAAGGCTCTAGAATTTGTAAAATACATAAATTCAAAGCTCCCTGGACTG  
CTAGAGCTTGAATATGAAGGGTTTTATAAGAGGGGATTCTTCGTTACGAAGAAGAGGTATGCAGTAATAGATGAA  
GAAGGAAAAGTCATTACTCGTGGTTTTAGAGATAGTTAGGAGAGATTGGAGTGAAATTGCAAAAGAAAACCAAGCT  
AGAGTTTTGGAGACAATACTAAAACACGGAGATGTTGAAGAAGCTGTGAGAATAGTAAAAGAAGTAATACAAAAG  
CTTGCCAATTATGAAATTCCACCAGAGAAGCTCGCAATATATGAGCAGATAACAAGACCATTACATGAGTATAAG  
GCGATAGGTCCTCACGTAGCTGTTGCAAGAACTAGCTGCTAAAGGAGTTAAAATAAAGCCAGGAATGGTAATT  
GGATACATAGTACTTAGAGGCGATGGTCCAATTAGCAATAGGGCAATTCTAGCTGAGGAATACGATCCCAAAAAG  
CACAAGTATGACGCAGAATATTACATTGAGAACCAGGTTCTTCAGCGGTACTTAGGATATTGGAGGGATTGGA  
TACAGAAAGGAAGACCTCAGATACCAAAAGACAAGACAAGTCGGCCTAACTTCTGGCTTAACATTAAAAAATCC

(SEQ ID NO.11)

It will be appreciated that preferred mutants encoded by the DNA sequences listed above comprise a codon for an alanine (A) inserted at position 2 that is not found in the wild type. Accordingly such mutants may be designated Y8A, Y38A and P116Δ mutants of the MAILDVDY form of Pfu Pol and correspond to Y7A, Y37A and P115Δ mutants of the true wild-type (MILDVDY).

It will also be appreciated that DNA sequences for other preferred mutants will be readily apparent given the abovementioned amino acid sequences. Accordingly DNA

molecules encoding Y7A, Y37A, V93A, I114R, I114Q and P115 $\Delta$  mutants of the true wild-type (MILDVDY) are also preferred nucleic acids according to the invention.

The variant polymerases as defined above are particularly useful for PCRs since they are thermally stable, have proofreading ability but are not stalled by the presence of dUTP.

Accordingly, a further aspect of the present invention provides a kit useful for polymerase chain reactions comprising DNA to be amplified, free bases, primers and a variant archaeal DNA polymerase having a modified amino acid sequence of a wild-type amino acid sequence, the modified sequence being in the amino-terminal amino acids that comprise a uracil binding pocket in the wild-type polymerase whereby the variant polymerase has reduced affinity for uracil compared to the wild-type polymerase.

The present invention further provides a method of amplifying DNA comprising the steps of (i) denaturing a double strand of DNA by heating a solution containing the DNA, free oligonucleotides, primers and a variant archaeal DNA polymerase having a modified amino acid sequence of a wild-type amino acid sequence, the modified sequence being in the amino-terminal amino acids that comprise a uracil binding pocket in the wild type polymerase whereby the variant polymerase has reduced affinity for uracil compared to the wild-type polymerase; (ii) reducing the temperature of the solution to effect annealing of the primer and the DNA and (iii) heating the solution to effect extension of DNA by the variant polymerase.

The ability of the variant DNA polymerase to amplify DNA in the presence of dUTP results in it being particularly suitable for PCR that uses dUTP rather than dTTP, for example in the prevention of contamination of samples.

A preferred protocol for carrying out PCR utilising variant polymerases according to the method of the present invention is as follows: PCR may be carried out under the following conditions: 100 µl volume, 20 mM Tris-HCl pH 8.8, 10 mM KCl, 10 mM (NH<sub>4</sub>)<sub>2</sub> SO<sub>4</sub>, 2 mM MgSO<sub>4</sub>, 0.1% Triton X100, 100 µg/ml BSA, 250 µM each of dATP, dGTP, dCTP and either 250 µM dTTP or 250 µM dUTP, 2.5 units DNA polymerase overlayed with 40 µl of mineral oil (1 unit of polymerase is defined as amount of enzyme that incorporates 10 nmol of dATP into acid-precipitable material using an activated calf-thymus DNA-based assay (4) in 30 min at 72°C). 5 ng of template DNA to be amplified is used and the concentration of the forward and reverse primers (each 18 bases in length) is 0.3 µM. Each PCR consisted of 30 cycles of 1 min at 95°C, 2 min at 52°C and 4.5 min at 72°C.

The present invention will now be further illustrated by means of the following Examples in which Example 1 investigates the uracil-binding pocket of Archaea polymerases, in particular the hyperthermophilic archaea, *Pyrococcus furiosus* (Pfu-pol) and Example 2 investigates the effect of mutagenesis of residues in and around the uracil-binding pocket of archaeal DNA polymerases, and with reference to the accompanying drawings in which:-

**Figure 1** illustrates an amino acid sequence alignment of archaeal family B DNA polymerases, corresponding to residues 1-130 of *Pyrococcus furiosus* polymerase. Candidates were identified using a WUBLAST search (European Bioinformatics Institute, [http://www.ebi.ac.uk/ebi\\_home.html](http://www.ebi.ac.uk/ebi_home.html)) for homologues of Pfu-Pol. An additional ENTREZ search of the SWISSPROT database for family B DNA polymerises was performed [Genbank (<http://www.ncbi.nlm.nih.gov/>)]. Sequence alignments were generated using ClustalX (version 1.81) [J. D. Thompson *et al.*, *Nucl. Acids. Res* **24**, 4876 (1997)]. The organisms and the DNA polymerase sequence accession numbers were: *Pyrococcus fuiosus* (Pfu) (P80061), *Thermococcus gorgonarius* (Tgo) (pdb 1D5A), *Pyrococcus kodakaraensis* (PKOD) (gi/13399597), *Desulfurococcus* strain Tok (DTok), *Thermococcus* sp. 9°N-7 (9°N-7) (Q56366), *Thermococcus litoralis* (Tli) (AAA72101.1), *Methanococcus voltae* (Mvo) (P52025), *Pyrobaculum islandicum* (Pis) (AAF27815.1), *Archaeoglobus fulgidus* (Agu) (O29753), *Cenarchaeaum symbiosum* (Csy) (AAC62712.1), *Sulfolobus acidocaldarius* (Sac) (P95690), *Sulfurisphaera ohwakuensis* (Soh) (BAA23994.1), *Sulfolobus solfataricus* (Sso) (P26811), *Pyrodictium occultum* (Poc) (BAA07579.1) and *Aeropyrum pernix* (Ape) (NP\_148473.1);

**Figure 2A:** The template-binding cleft T (21) of Tgo-Pol showing the presence of a pocket. **B:** The N-terminal domain of Tgo-Pol with amino acids that form the pocket shown in space-fill: Y7; P36/Y37; amino acids 90-97 (22); amino acids 112-116 (23).  $\alpha$ -Helices are shown (24) and  $\beta$ -sheets (25). **C:** Amino acid sequences of the N-terminal domains of Tgo-Pol (upper sequence (SEQ ID NO.12)) and RB69-Pol (lower sequence (SEQ ID NO.13)). Amino acids that form the pocket in Tgo-Pol (and the corresponding residues in RB69-Pol)

are underlined and correspond to the amino acids identified in panels B and E. Cylinders represent  $\alpha$ -helices and arrows  $\beta$ -sheets. The amino acid sequences have minimal homology and have been aligned using structural homology. **D:** Structural alignment of the N-terminal domains of Tgo-Pol and RB69-Pol. The insert in Tgo-Pol is shown (26). This structural superimposition was used to generate the amino acid alignment shown in C. **E:** The N-terminal domain of RB69-Pol. Space-filled amino acids (V8 Q10; residues 65-72 (27); residues 84-89 (28); P35/S36 lie substantially behind residues 84-89) correspond to those in Tgo-Pol (shown in panel B) which form the pocket. Both V8 and Q10 are near the position of the Tgo-Pol Y7.  $\alpha$ -Helices are shown (24) and  $\beta$ -sheets (25). Images/structural homology models were produced using Swiss-Model [N. Guex, M. C. Peitsch, *Electrophoresis* 18, 2714 (1997)] (<http://www.expasy.ch/spdbv/>), POV-Ray [C. Cason, POV-Ray for Windows, version 3.1g (1999)] (<http://www.povray.org>) and Rasmol [R. Sayle, J. F. Milner-White, *Trends Biochem. Sci.* 20, 374 (1995)].

**Figure 3:** Modelling pyrimidines into the N-terminal domain pocket of Tgo-Pol using Web Lab Viewer Pro [Molecular Simulations Inc., Web Lab Viewer Pro (version 4.0) (2000)] (<http://www.msi.com>). Hatched lines represent hydrogen bonds; except steric clashes, are identified (31). Uracil formed four enzyme-base hydrogen bonds and no clashes. Cytosine and thymine resulted in fewer hydrogen bonds and/or clashes.

**Figure 4 A and B:** Primer extension reactions using a 5'-<sup>32</sup>P labeled 24-mer primer (5'-GGGGATCCTCTAGAGTCGACCTGC-3' (SEQ ID NO.14)) (2.5nM)

annealed to 44-mer template (5'-GGAGACAAGCTTG(U/T)ATGCCTGCAGGTCGACTCTAGCGGCTAAAA-3' (SEQ ID NO.15)) (5nM). The primer hybridises with the underlined section of the template to place a uracil (thymine in controls) seven bases from the primer-template junction (H.H. Hogrefe, C. J. Hansen, B. R. Scott, K. B. Nielson, *Proc. Natl. Acad. Sci. U.S.A.*, **99**, 596 (2002)). In the presence of the four dNTPs all Pfu-Pol variants were able to fully copy the control template, lacking uracil, converting the labeled 24-mer primer into a product 44 bases long (A). With uracil in the template the wild-type enzyme (WT) stalled polymerisation (H.H. Hogrefe, C. J. Hansen, B. R. Scott, K. B. Nielson, *Proc. Natl. Acad. Sci. U.S.A.*, **99**, 596 (2002)) giving a truncated product (B). The arrow (41) indicates the major pause site, shown (using standards, not illustrated) to occur four base before uracil. With Y7A, V93Q and P115Δ some full-length product was observed (amounts: V93Q > P115Δ > Y7A) indicating read-through of the template-strand uracil (panel B). No full-length product was seen with Y37A, Y37F and P115F (some material partially extended past the "uracil induced pause" was seen with Y7A, Y37A and Y37F. The first and second lanes contain standard 24-mer and 44-mer. **C and D:** PCR reactions. In control PCR with the four normal dNTPs (dATP, dGTP, dCTP and dTTP) all the Pfu-Pol variants (and Taq-Pol) gave the anticipated PCR product, ~ 2 kbases in length (C). When dUTP replaced dTTP, Taq-Pol produced a 2 kbase fragment (D). With most Pfu-Pol variants (WT, Y37A, Y37F and P115F) no PCR product was seen (D). Three Pfu-Pol mutant's produced a PCR product (amounts: V93Q > P115Δ > Y7A) (2 x and 4 x loadings, respectively, were used for visualisation with P115Δ and Y7A). All PCR reactions were carried out under identical conditions with no optimisation; this probably accounts for the



lower weight contaminant seen in some lanes. Markers (important sizes indicated) were from Promega.

**Figure 5:** Polymerase chain reaction (PCR) with *Thermus aquaticus* polymerase (Taq-Pol) and *Pyrococcus furiosus* polymerase (Pfu-Pol) (wild type and two mutants, V93Q and V93R). A DNA fragment (~ 1 kilobase in length) from the plasmid pET-17b(Pfu-Pol) was amplified. Conditions 20 mM Tris-HCl, pH 8.8, 10 mM KCl, 10 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 2 mM MgSO<sub>4</sub>, 0.1 % (v/v) Triton X100 and 0.1 mg/ml bovine serum albumin. Each reaction contained 250  $\mu$ M each dATP, dGTP and dCTP. Reactions 1-4 contained 250  $\mu$ M TTP; reactions 5-8 250  $\mu$ M dUTP. 2.5 units of the polymerase were used. Cycle 1 x 10 minutes 94 °C; 30 x 1minute 94 °C/2 minutes 42°C/1 minute 72 °C; 1 x 10 minutes 72 °C. The Pfu-Pol mutants V93Q and V93R give higher yield of PCR product (lanes 3,4) than wild type Pfu-Pol (lane 2) when TTP is used. Wild type Pfu-Pol does not give a PCR product with dUTP (lane 6), whereas Pfu-Pol mutants V93Q and V93R give a product (lanes 7,8) [lane 1= Taq-Pol with TTP; lane 2= Wild type Pfu-pol with TTP; lane 3= Pfu-Pol V93Q with TTP; lane 4= Pfu-Pol V93R with TTP; lane 5= Taq-Pol with dUTP; lane 6= Wild type Pfu-pol with dUTP; lane 7= Pfu-Pol V93Q with dUTP; lane 8= Pfu-Pol V93R with dUTP] [M= 100bp ladder; Mb= 1Kb ladder].

### **Example 1**

Crystal structures are known for five family B DNA polymerases; one viral, the remaining four archaeal. The first structure to be solved was for the bacteriophage RB69 polymerase (RB69-Pol) (J. Wang *et al.*, *Cell* 89, 1087 (1997)); a structure with primer-template has also been determined (M. C. Franklin, J. J. Wang, T. A. Steitz, *Cell* 105, 657 (2001)). More recently, the structure of an archaeal family B DNA polymerase, from the hyperthermophilic archaeon *Thermococcus gorgonarius* (Tgo-Pol), has been reported (K. P. Hopfner *et al.*, *Structure* 7, 1189 (1999)). Three other archaeal polymerase structures, *Desulfurococcus* strain Tok (DTok-Pol) (Y. Zhao *et al.*, *Structure* 7, 1189 (1999)), *Thermococcus* sp. 9°N-7 (9°N-7-Pol) (A. C. Rodriguez, H-W. Park, C. Mao, L. S. Beese, *J. Mol. Biol.* 299, 447 (2000)) and *Pyrococcus kodakaraensis* KOD1 (KOD1-Pol) (H. Hashimoto *et al.*, *J. Mol. Biol.* 306, 469 (2001)), were subsequently solved. Only apo-enzyme structures are known for the archaea. All five family B polymerases contain five distinct domains, the N-terminal domain, the exonuclease or 'editing' domain and three polymerase active site domains. The folding of the five domains forms three distinct clefts extending from a central hole. Two (named clefts D and T) are oriented approximately 180° relative to each other, on either side of the central hole. The structure of RB69-pol containing a primer-template demonstrates that cleft D binds double stranded primer-template and cleft T binds single-stranded template (M. C. Franklin *et al supra*). The three polymerase domains forms cleft D, whereas cleft T is formed by the exonuclease domain and the N-terminal domain. The third cleft is perpendicular to the other two and represents the 3'-5' exonuclease/editing cleft.

Examination of cleft T in the case of Tgo-Pol (the other archaeal polymerases give similar results) revealed the presence of a pocket located on a surface exposed face towards the outer edge of the polymerase (figure 2A). The location of this pocket, in the template strand binding region, approximately four bases from the primer-template junction makes it a clear candidate for uracil recognition. The pocket is comprised of amino acids that are solely present in the N-terminal domain. Amino acids from four regions of this domain, which are close together in space, are used to assemble the putative uracil binding pocket. These amino acids are illustrated in figure 2B (a structural representation of the N-terminal domain) and underlined in figure 2C (the amino acid sequence of the N-terminal domain). The function of Y7, which sits at the entrance of the pocket, is obscure. It may form a lid, which closes following binding, ensuring trapping (and hence high affinity) of uracil. The base of the pocket (clearly visible as P36/Y37 in figure 2B) is formed by Y37, oriented by P36 at the beginning of a  $\beta$  sheet and supported by K84 "underneath" Y37. One side of the pocket is formed by amino acids 90-97, present in a  $\alpha$ -helix. A proline (P94) bends the  $\alpha$ -helix, forming a curved wall (figure 2B). The other side comprises residues 110-116; amino acids present in a loop region (110-114) or at the beginning of a second  $\alpha$ -helix (115-116). This  $\alpha$ -helix commences with amino acids P115 and F116. The proline appears to have a critical role, ensuring that P115 and F116 are able to form part of the pocket's curved wall (figure 2B).

The viral polymerase from RB-69 shows 61% amino acid identity to bacteriophage T4-Pol. Additionally, the N-terminal of RB69-Pol is identical to a structure for an N-terminal fragment of T4-Pol (J. Wang, P. Yu, T. C. Lin, W. H. Konigsberg, T.A. Steitz, *Biochemistry* 35, 8110 (1996)). Previously, it has been demonstrated that T4-Pol did not stall

polymerisation in response to template-strand uracil (M. A. Greagg *et al.*, *Proc Natl. Acad. Sci. U.S.A.* 96, (1999)); inability to recognize uracil would also be expected for RB69-Pol. This contrasts with Tgo-Pol which stalls polymerization when template strand uracil is encountered (M. A. Greagg *et al.*, *Supra*). Tgo-Pol is also inhibited by uracil containing DNA (R. S. Lasken, D. M. Schuster, A. Rashtchian, *J. Biol. Chem.* 271, 17692 (1996)); a feature not seen with viral enzymes. Therefore, if the pocket seen with Tgo-Pol is used for uracil detection, it should be absent for the viral enzymes. The N-terminal domains of Tgo-Pol and RB69-Pol show considerable structural homology (figure 2D); near perfect alignment of secondary structural elements is observed, despite the almost complete lack of amino acid sequence homology (figure 2C). The only significant difference is the presence of an amino acid insert (shown in figure 2D) in the archaeal enzyme. However, detailed comparisons demonstrate differences in the two polymerases (figure 2E). With Tgo-Pol, Y7 may act as a lid and Y36 forms the base of the pocket; in RB-69 these residues are replaced with valine/glutamine (both these amino acids in RB69-Pol are near Y7 of Tgo-Pol and it is not clear which is the exact replacement) and serine respectively. Similarly Tgo-Pol uses a proline-containing kinked  $\alpha$ -helix (residues 90-97) to form one wall of the pocket. In RB69-Pol the corresponding  $\alpha$ -helix (residues 65-74) lacks the proline; therefore, the helix is straight and does not form a good wall. However, a key feature involves P115 and F116, which form part of one of the walls of the pocket with Tgo-Pol. In the case of RB69 the proline is missing and this results in the corresponding phenylalanine (F88) falling into and completely filling the pocket. As shown in figure 1E this means that the viral enzyme lacks a uracil binding pocket (P35/S36 are substantially obscured). The differences between the viral

and archaeal enzymes, based on subtle changes to a few amino acids, provide compelling evidence that the N-terminal pocket is responsible for uracil detection.

It was possible to model uracil into the pocket of Tgo-Pol (figure 3A). The most favourable orientation produces four hydrogen bonds between the protein and uracil. In all cases the protein uses the peptide backbone for hydrogen bond formation. The interactions comprise: I114 (peptide -NH) to uracil C2 =O group; E111 (peptide =O) to uracil N3H; Y37 (peptide =O) to uracil N3H and (peptide -NH) to uracil C4=O (figure 3A). Cytosine superimposed at the same position as uracil forms only one hydrogen bond (I114 (peptide -NH) to uracil C2 =O) and the 4-NH<sub>2</sub> group clashes with the main chain atoms of Y37 (peptide =O and -NH) (figure 3B). This clash could be relieved by repositioning the cytosine, but only at the expense of the one H-bond, resulting in no interactions between the base and the protein. Thymine superimposed at the uracil position could form most of the protein-base hydrogen bonds (interactions were identical to uracil except the Y37 (peptide -NH) to uracil/thymine N3H hydrogen bond was not formed). Critically the C5 -CH<sub>3</sub> group showed a severe steric clash with the edges of the cyclic P36 side chain and the ring of F116, thereby preventing binding of the base within the pocket (figure 3C). Thus the pocket is highly specific for binding uracil and able to discriminate against the 'normal' DNA pyrimidines.

An amino acid sequence alignment has been carried out for the N-terminal domains of fourteen archaeal family B DNA polymerases (figure 1); eight were from the crenarchaea and six from the euryarchaea. Twelve of the polymerases were either thermophiles or hyperthermophiles, one was mesophilic (*Methanococcus voltae* (Mvo)) and one was psychrophilic (*Cenarchaeum symbiosum* (Csy)). Two highly conserved regions (A and B) which contain most of the amino acids that form the uracil binding pocket, are seen. Many of

the amino acids comprising the uracil binding pocket (figure 2B and 2C) are highly conserved; especially uracil-contacting residues (figure 3A). Thus P36, Y37, E111 and I114 show 100 % identity. The possible pocket lid, Y7, also shows 100 % conservation. Several other key amino acids e.g. V93 (which lines one side of the pocket), P115 and F116 (which line the other side of the pocket) show a high degree of conservation.

### Example 2

Use was made of three assays to test the ability of the mutant polymerases to recognise uracil. Primer extension reactions (M. A. Greagg *et al.*, *Supra*) measure the ability of a polymerase to extend a primer through uracil bases in the template strand. As expected both the wild type and the mutant enzymes were able to completely copy a control template, lacking uracil (figure 4A). As previously observed (M. A. Greagg *et al.*, *Supra*), the wild type enzyme stalled polymerisation four bases upstream of template-strand uracil, resulting in a truncated product (figure 4B). Y37A, Y37F and P115F behaved in a similar manner to wild type. However, three of the mutant enzymes, V93Q, P115Δ and Y7A, produced full-length product when uracil was present (figure 4B). With V93Q full-length product predominated; in the cases of P115Δ and Y7A both full-length and truncated product were seen.

Next the ability of the polymerases to bind single stranded DNA containing uracil was investigated, using a binding assay based on fluorescence anisotropy (S. L. Reid, D. Parry, H.-H. Liu, B. A. Connolly, *Biochemistry* 40, 2484 (2001)).  $K_D$  was determined by fluorescence anisotropy used an oligodeoxynucleotide containing a single uracil and a hexachlorofluorescein label at its 5'-terminal. The oligodeoxynucleotide used was: 5'-hex-GCCCGCGGGAUATCGGCCCTTA-3' (SEQ ID NO.16) (or a control in which the uracil was replaced with thymine). The concentration of the oligodeoxynucleotide was 5 nM in 1 ml

of 10 mM Hepes-NaOH, pH7.5, 100mM NaCl, 1mM EDTA. Aliquots of the enzyme were added and the anisotropy measured; titration was continued until the anisotropy stopped increasing. Data fitting to obtain  $K_D$  values is as described by Reid *et al* (*Supra*).  $K_D$  values are summarised in table 1. The wild type enzyme bound the uracil-containing oligodeoxynucleotide with a  $K_D$  of 8.3 nM, a 17-fold preference over a control strand lacking this base. Three of the mutants, Y7A, V93Q and P115 $\Delta$  bound to the uracil-containing oligodeoxynucleotide less well than the wild type ( $K_D$  values of 25.7, 144.5 and 84 nM respectively, table 1). These mutants correspond exactly to those able to read through a uracil-containing template; furthermore the diminution in uracil binding corresponds with read through capability. In both assays loss of uracil recognition is: V93Q > P115 $\Delta$  > Y7A. These three mutants also show a reduced preference for uracil-containing DNA over the control sequence; with P115 $\Delta$  and V93Q the preference is virtually abolished. The mutants Y7A, Y37A and P115F bind uracil-containing DNA with essentially the same affinity as the wild type (table 1). In some cases, Y37A and Y37F, the preference for the uracil-containing oligodeoxynucleotide is reduced, but this arises solely from tighter binding of the control. Table 1 also gives the specific activity of the mutant Pfu-Pols relative to the wild type. In general there are only small decreases in activity with even the mutant with the lowest activity (Y37A) retaining 38 % of the wild type activity.

The DNA polymerase activity assay (Richardson, C. C. (1966) in *Procedures in Nucleic Acids Research*, G. L. Cantoni, D. R. Davies, Eds, (Harper and Row, New York, 1966), pp. 263-27) used 50  $\mu$ l samples containing 20 mM Tris-HCl, pH 8.8, 10mM KCl, 10mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 2 mM MgSO<sub>4</sub>, 0.1% Triton X100, 100 $\mu$ g/ml BSA, 200 $\mu$ M each dNTPs, 0.2mg/ml activated calf thymus DNA (AP Biotech), 1 $\mu$ Ci 3000Ci/mmol [ $\alpha$ -<sup>32</sup>P] dATP. Pfu-

Pol (the amount varied depending on the activity of the enzyme) was added and a 10 minute incubation at 72°C was used (reactions were linear over this time). After this period the amount of radioactivity incorporated into acid-precipitable material was determined by scintillation counting. 1 unit of enzyme is defined as amount of enzyme that incorporates 10 nmol of dATP into acid-precipitable material in 30 min at 72°C.

**Table 1.** Pfu-Pol variants: specific activity and ability to bind to uracil containing DNA.

Pfu-Pol variant	specific activity (units/mg)	% activity (relative to wild type)	K <sub>D</sub> (nM) (uracil)	K <sub>D</sub> (nM) (control)	Preference for uracil
wild-type	3556	100	8 ± 1	140 ± 10	18
Y7A	2314	65	26 ± 1	255 ± 20	10
Y37A	1352	38	9 ± 2	57 ± 7	6
Y37F	2169	61	10 ± 4	120 ± 13	12
V93Q	1643	46	144 ± 7	277 ± 22	2
P115F	2541	71	7 ± 1	148 ± 10	21
P115Δ	1637	46	84 ± 4	111 ± 5	1.3

The specific activities of the Pfu-Pol variants was determined using the incorporation of [ $\alpha$ -<sup>32</sup>P]-dATP into acid-precipitable calf-thymus DNA as described (see above). Values are accurate ± 15 %. Binding constants were determined by fluorescence anisotropy using hex-GCCCGCGGGAUATCGGCCCTTA (SEQ ID NO.16) (uracil) or an analogous oligodeoxynucleotide in which the uracil was replaced with thymine (control) (S. L. Reid, *et al*; *supra* and J. Wang *et al.*, (*Supra*)). Each value was determined three times and the average ± one standard deviation is given. The preference for uracil is the ratio  $K_D(\text{uracil})/K_D(\text{control})$ .



Finally PCR was performed +/- dUTP the ability of Pfu-Pol to carry out PCR was evaluated by amplifying a – 2 kbase fragment (between the T7 promoter and the *Hind*III site) of pET17-b(Pfu-Pol) [S. J. Evans *et al.*, *Nucl. Acids. Res.* 28, 1059 (2000)]. Conditions: 100µl volume, 20 mM Tris-HCl pH 8.8, 10 mM KCl, 10 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 0.1% Triton X100, 100 µg/ml BSA, 250 µM each dNTP (one set of reactions contained dTTP, the other dUTP), 2.5 units DNA polymerase overlayed with 40 µl of mineral oil. 5 ng of pET17-b(Pfu-Pol) was used and the concentrations of the forward and reverse primers were both 0.3 µM. Each reaction was 30 cycles of 1 min at 95°C, 2 min at 52°C and 4.5 min at 72°C. PCR with Taq-Pol was identical to Pfu-Pol save 10 mM Tris-HCl pH 8.8, 50 mM KCl, 0.08% NP-40, 1.5 mM MgCl<sub>2</sub> was used. Analysis used ethidium bromide-stained agarose gels.

Pfu-Pol (wild type and mutants) and Taq-Pol, were able to perform PCR with the four normal dNTPs (figure 4C). When dUTP was used in place of dTTP, PCR with Taq-Pol was unaffected; however, wild-type Pfu-Pol gave no product (figure 4D). The three mutants that showed diminished uracil recognition in read-through and binding assays, Y7A, P115Δ and V93Q, gave a PCR product (figure 4D). The amount of PCR product produced was V93Q > P115Δ > Y7A, again matching the order found for loss of uracil recognition. dUTP (concentration 250 µM) completely replaces dTTP in these reactions. The wild type enzyme is completely inhibited when 0.02 µM dUTP is used to spike PCR reactions containing the four normal dNTPs (H. H. Hogrefe, *et al* (*Supra*)); clearly showing these three mutants are very disabled in uracil recognition. The Pfu-Pol mutants, Y37A, Y37F and P115F did not give a PCR product with dUTP.

Figure 5 shows the results of PCR with *Thermus aquaticus* polymerase (Taq-Pol), the wild type and two mutants (V93Q and V93R) of *Pyrococcus furiosus* polymerase (Pfu-Pol). PCR amplification was performed under two distinct sets of conditions, i.e. in the presence of TTP and in the presence of dUTP.

As expected, all four polymerases, Taq-Pol, the wild type of Pfu-pol and the two mutations of Pfu-pol (V93Q, V93R), were able to successfully mediate amplification of the DNA sample in the presence of TTP. Clearly visible bands of 1064 bp fragments in lanes 1 to 4 illustrate this successful amplification. The use of dUTP resulted in the expected amount of amplification when used together with Taq-Pol as illustrated in lane 5. Moreover, lane 6 shows no amplified DNA sample and thereby confirmed that dUTP induces blockage of Pfu-Pol mediated amplification. Most importantly, however, lanes 7 and 8 show bands corresponding to 1064 bp fragments and thus confirm that both Pfu polymerase mutants are capable of amplification. This proves that, unlike wild type Pfu-Pol, the V93Q and V93R mutants of Pfu-Pol are not affected by dUTP-induced blockage.

Overall, Figure 5 confirms the usefulness of the mutants according to the invention in PCR utilizing dUTP.

### CLAIMS

1. A variant archaeal DNA polymerase having a modified amino acid sequence of a wild-type amino acid sequence, the modified sequence being in the amino-terminal amino acids that comprise a uracil-binding pocket in the wild-type polymerase whereby the variant polymerase has reduced affinity for uracil than the wild-type polymerase.
2. A variant archaeal DNA polymerase according to claim 1 having a modified amino acid sequence of wildtype polymerases selected from *Thermococcus gorgonarius* (Tgo-Pol), *Thermococcus litoralis* (Tli-Pol), *Thermococcus* sp. 9°N-7 (9°N-7-Pol), *Desulfurococcus* strain Tok (DTok-Pol), *Pyrobaculum islandicum* (Pis-Pol), *Archaeoglobus fulgidus* (Afu-Pol), *Sulfolobus acidocaldarius* (Sac-Pol), *Sulfurisphaera ohwakuensis* (Soh-Pol), *Sulfolobus solfataricus* (Sso-Pol), *Pyrodictium occultum* (Poc-Pol) or *Aeropyrum pernix* (Ape-Pol).
3. A variant archaeal DNA polymerase according to claim 1 having a modified amino acid sequence of wildtype *Pyrococcus furiosus* DNA polymerase (Pfu-Pol),
4. A variant archaeal DNA polymerase according to claim 3 having modifications in amino acids 1-40 or amino acids 78-130.
5. A variant archaeal DNA polymerase according to claim 4 of SEQ ID NO.2 having modifications to amino acids 7, 36, 37, 90-97 or 112 - 119.

6. A variant archael DNA polymerase according to claim 5 having modifications to amino acids Y7, Y37, V93, I114 or P115.
7. A variant archaeal DNA polymerase according to claim 5 wherein the modification is Y7A.
8. A variant archaeal DNA polymerase according to claim 5 wherein the modification is Y37A.
9. A variant archaeal DNA polymerase according to claim 5 wherein the modification is V93Q.
10. A variant archaeal DNA polymerase according to claim 5 wherein the modification is V93R.
11. A variant archaeal DNA polymerase according to claim 5 wherein the modification is I114R.
12. A variant archaeal DNA polymerase according to claim 5 wherein the modification is I114Q.
13. A variant archaeal DNA polymerase according to claim 5 wherein the modification is P115A.

14. A variant archaeal DNA polymerase according to claim 4 of SEQ ID NO.1 having modifications to amino acids 8, 37, 38, 91-98 or 113 – 120.
15. A variant archaeal DNA polymerase according to claim 14 having modifications to amino acids Y8, Y38, V94, I115 or P116.
16. A variant archaeal DNA polymerase according to claim 14 wherein the modification is Y8A.
17. A variant archaeal DNA polymerase according to claim 14 wherein the modification is Y38A.
18. A variant archaeal DNA polymerase according to claim 14 wherein the modification is V94Q.
19. A variant archaeal DNA polymerase according to claim 14 wherein the modification is V94R.
20. A variant archaeal DNA polymerase according to claim 14 wherein the modification is I115R.
21. A variant archaeal DNA polymerase according to claim 14 wherein the modification is I115Q.

22. A variant archaeal DNA polymerase according to claim 14 wherein the modification is P116Δ.
23. A variant archaeal DNA polymerase according to any preceding claim having modifications in the amino acid motif: E - - I -F/Y- - -Y- -D.
24. A nucleic acid molecule encoding an archaeal DNA polymerase according to any one of claims 1 - 23.
25. A method of amplifying DNA comprising the steps of (i) denaturing a double strand of DNA by heating a solution containing the DNA, free oligonucleotides, primers and a variant archaeal DNA polymerase as defined in any one of claims 1 - 23; (ii) reducing the temperature of the solution to effect annealing of the primer and the DNA and (iii) heating the solution to effect extension of DNA by the variant polymerase.
26. A kit useful for polymerase chain reactions comprising a variant archaeal DNA polymerase as defined in any one of claims 1 - 23 and optionally DNA to be amplified, free bases and primers.

Pfu	1	----	MI	EDTD	MTTE	EXP	WIR	LEKK	----	EN	GE	KI	H	D	R	E	R	----	MI	YA	40
Tgo	1	----	MI	EDTD	MTTE	EXP	WIR	LEKK	----	EN	GE	KI	I	D	R	E	R	----	MI	YA	40
PKOB	1	----	MI	EDTD	MTTE	EXP	WIR	LEKK	----	EN	GE	KI	E	D	R	T	E	----	ET	YA	40
UTok	1	----	MI	EDTD	MTTE	EXP	WIR	LEKK	----	EN	GE	KI	I	D	R	E	R	----	MI	YA	40
9°N-7	1	----	MI	EDTD	MTTE	EXP	WIR	LEKK	----	EN	GE	KI	E	D	R	T	E	----	ET	YA	40
Tli	1	----	MI	EDTD	MTTE	EXP	WIR	LEKK	----	EN	GE	KI	E	D	R	T	E	----	ET	YA	40
Nvo	1	----	MI	EDTD	MTTE	EXP	WIR	LEKK	----	EN	GE	KI	E	D	R	T	E	----	ET	YA	40
Pis	1	----	MI	EDTD	MTTE	EXP	WIR	LEKK	----	EN	GE	KI	E	D	R	T	E	----	ET	YA	40
Afu	1	----	MI	EDTD	MTTE	EXP	WIR	LEKK	----	EN	GE	KI	E	D	R	T	E	----	ET	YA	40
Csy	2	----	MI	EDTD	MTTE	EXP	WIR	LEKK	----	EN	GE	KI	E	D	R	T	E	----	ET	YA	40
Sac	40	----	MI	EDTD	MTTE	EXP	WIR	LEKK	----	EN	GE	KI	E	D	R	T	E	----	ET	YA	40
Soh	41	----	MI	EDTD	MTTE	EXP	WIR	LEKK	----	EN	GE	KI	E	D	R	T	E	----	ET	YA	40
Sso	41	----	MI	EDTD	MTTE	EXP	WIR	LEKK	----	EN	GE	KI	E	D	R	T	E	----	ET	YA	40
Poc	77	----	MI	EDTD	MTTE	EXP	WIR	LEKK	----	EN	GE	KI	E	D	R	T	E	----	ET	YA	40
Rpe	121	----	MI	EDTD	MTTE	EXP	WIR	LEKK	----	EN	GE	KI	E	D	R	T	E	----	ET	YA	40

1

2

FIG.1





																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																															</
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	----






FIG.1 continued

**FIG. 2**



**FIG.3**





FIG. 4

FIG. 5

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 03/01623

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C12N9/12 C12N15/03

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched.

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data, EMBASE, BIOSIS, SEQUENCE SEARCH, MEDLINE

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	FOGG MARK J ET AL: Structural basis for uracil recognition by archaeal family B DNA polymerases." NATURE STRUCTURAL BIOLOGY. UNITED STATES DEC 2002, vol. 9, no. 12, December 2002 (2002-12), pages 922-927, XP00210141 ISSN: 1072-8368 the whole document  --- -/--	1-15, 23-26

☒ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex.

## \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the International filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the International filing date but later than the priority date claimed

\*T\* later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*Z\* document member of the same patent family

Date of the actual completion of the international search

5 August 2003

Date of mailing of the International search report

22/08/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Armandola, E

## INTERNATIONAL SEARCH REPORT

Int: International Application No  
PCT/GB 03/01623

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	GREAGG M A ET AL: "A read-ahead function in archaeal DNA polymerases detects promutagenic template-strand uracil." PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA. UNITED STATES 3 AUG 1999, vol. 96, no. 16, 3 August 1999 (1999-08-03), pages 9045-9050, XP002250142 ISSN: 0027-8424	
A	EVANS S J ET AL: "Improving dideoxynucleotide-triphosphate utilisation by the hyper-thermophilic DNA polymerase from the archaeon Pyrococcus furiosus." NUCLEIC ACIDS RESEARCH. ENGLAND 1 MAR 2000, vol. 28, no. 5, 1 March 2000 (2000-03-01), pages 1059-1066, XP002164138 ISSN: 1362-4962	
T	CONNOLLY B A ET AL: "Uracil recognition by archaeal family B DNA polymerases." BIOCHEMICAL SOCIETY TRANSACTIONS. ENGLAND JUN 2003, vol. 31, no. Pt 3, June 2003 (2003-06), pages 699-702, XP009015138 ISSN: 0300-5127	

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**